



ELSEVIER

Contents lists available at ScienceDirect

## Heart &amp; Lung

journal homepage: [www.heartandlung.com](http://www.heartandlung.com)

## Corrigendum

## Corrigendum to “The effect of Shiatsu massage on agitation in mechanically ventilated patients: A randomized controlled trial “[Heart & Lung Volume 50 (2021) 893-897]

Mehdi Harorani<sup>1,2,\*</sup>, Masoumeh Garshasbi<sup>3</sup>, Mohamad Sediqi<sup>3</sup>, Zohreh Farahani<sup>1</sup>,  
Danial Habibi<sup>4</sup>, Mahtab Farahani<sup>3</sup>, Nazanin Amini<sup>5</sup>, Zahra Velashjerdi<sup>6</sup>

<sup>1</sup> Department of Nursing, School of Nursing, Arak University of Medical Sciences, Arak, Iran

<sup>2</sup> Traditional and Complementary Medicine Research Center (TCMRC), Arak University of Medical Sciences, Arak, Iran

<sup>3</sup> Student Research Committee, Department of Nursing, School of Nursing, Arak University of Medical Sciences, Arak, Iran

<sup>4</sup> Department of Biostatistics and Epidemiology, Faculty of Health, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>5</sup> Department of paramedicine, Arak University of Medical Sciences, Arak, Iran

<sup>6</sup> Master of Nursing, Valiasr Arak Hospital, Arak University of Medical Sciences, Arak, Iran

The authors regret that the printed version of the above article contained a number of errors and miscalculations. In table number one of the published article, the number of samples, P-value and the number of some variables were typed incorrectly and must be corrected as follows.

The authors would like to apologise for any inconvenience caused.

**Table 1**  
Comparison of demographic characteristics in the control and intervention groups.

Variable		intervention group) N=33 No%	control group (N=34) No%	P-value
<b>Sex</b>	Male	17 (51.5)	17 (50)	0.901*
	Female	16 (48.5)	17 (50)	
<b>Educational level</b>	Illiterate	12 (36.4)	12 (35.3)	0.582**
	Diploma and Less than diploma	18 (54.5)	21 (61.8)	
	Collegiate	3 (9.1)	1 (2.9)	
<b>History of underlying disease</b>	yes	24 (72.7)	17 (50)	0.056**
	no	9 (27.3)	17 (50)	
<b>Marital status</b>	Single	3 (9.1)	7 (20.6)	0.130**
	Married	30 (90.9)	25 (73.5)	
	Widow	0	2 (5.9)	
	Divorced	0	0	
<b>Tranquilizer used during the intervention.</b>	Apotel	3 (9.1)	3 (8.9)	0.288*
	Pethidine	2 (6.1)	1 (2.9)	
	Fentanyl	5 (15.2)	12 (35.3)	
	Morphine sulfate	10 (30.3)	5 (14.7)	
	Not receiving sedation	13 (39.3)	13 (38.2)	
<b>Sedative used during the intervention.</b>	Midazolam	19 (57.6)	13 (38.2)	0.230*
	Propofol	1 (3)	1 (2.9)	
	Non-use	13 (39.4)	20 (58.9)	

All values are expressed as number (percentage) or mean ± standard deviation.

\* Chi-square

\*\* Fisher exact test.

DOI of original article: <http://dx.doi.org/10.1016/j.hrtlng.2021.07.013>.

\* Corresponding author: Mehdi Harorani, Department of Nursing, School of Nursing, Arak University of Medical Sciences, Arak, Iran  
E-mail address: [m.harorani@yahoo.com](mailto:m.harorani@yahoo.com) (M. Harorani).

<https://doi.org/10.1016/j.hrtlng.2023.02.008>

0147-9563/© 2021 Elsevier Inc. All rights reserved.